

After you get the COVID-19 vaccine, you still need to:



Wear a cloth mask or face covering when you are out in public.



Avoid close contact with people by staying 6 feet apart from others.



Wash your hands often with soap and water for at least 20 seconds.



Cover coughs and sneezes with your arm or inner elbow, and avoid touching your eyes, nose and mouth with your hands.



Stay home if you are sick or have symptoms of COVID-19.

Learn more about your COVID-19 vaccine on the reverse side. >>>

After you get the COVID-19 vaccine, you still need to:



Wear a cloth mask or face covering when you are out in public.



Avoid close contact with people by staying 6 feet apart from others.



Wash your hands often with soap and water for at least 20 seconds.



Cover coughs and sneezes with your arm or inner elbow, and avoid touching your eyes, nose and mouth with your hands.



Stay home if you are sick or have symptoms of COVID-19.

Learn more about your COVID-19 vaccine on the reverse side. >>>

You received the Pfizer vaccine.

- You need **2 doses** for maximum protection.
- Your **second dose appointment date and time** is written on the back of your vaccination record card.
- Please arrive at your scheduled appointment time.
- You will receive a text and/or email reminder for your second dose appointment 2 days prior to your scheduled date and time.
- Please **bring your vaccine record card** to your second appointment.



Common Side Effects *(usually more after second dose)*

- Injection site soreness
- Chills
- Fatigue
- Fever
- Headache
- Joint pain
- Muscle aches


Side effects typically last 2-3 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519



3/1/2021

You received the Pfizer vaccine.

- You need **2 doses** for maximum protection.
- Your **second dose appointment date and time** is written on the back of your vaccination record card.
- Please arrive at your scheduled appointment time.
- You will receive a text and/or email reminder for your second dose appointment 2 days prior to your scheduled date and time.
- Please **bring your vaccine record card** to your second appointment.



Common Side Effects *(usually more after second dose)*

- Injection site soreness
- Chills
- Fatigue
- Fever
- Headache
- Joint pain
- Muscle aches


Side effects typically last 2-3 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519



3/1/2021

You received the Moderna vaccine.

- You need **2 doses** for maximum protection.
- Your **second dose appointment date and time** is written on the back of your vaccination record card.
- Please arrive at your scheduled appointment time.
- You will receive a text and/or email reminder for your second dose appointment 2 days prior to your scheduled date and time.
- Please **bring your vaccine record card** to your second appointment.



Common Side Effects *(usually more after second dose)*

- Injection site soreness
- Chills
- Fatigue
- Swollen lymph nodes in same arm as injection
- Fever
- Headache
- Nausea
- Joint pain
- Muscle aches
- Vomiting


Side effects typically last 2-3 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519



You received the Moderna vaccine.

- You need **2 doses** for maximum protection.
- Your **second dose appointment date and time** is written on the back of your vaccination record card.
- Please arrive at your scheduled appointment time.
- You will receive a text and/or email reminder for your second dose appointment 2 days prior to your scheduled date and time.
- Please **bring your vaccine record card** to your second appointment.



Common Side Effects *(usually more after second dose)*

- Injection site soreness
- Chills
- Fatigue
- Swollen lymph nodes in same arm as injection
- Fever
- Headache
- Nausea
- Joint pain
- Muscle aches
- Vomiting


Side effects typically last 2-3 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519



You received the Johnson & Johnson vaccine.

- You need **1 dose** for this vaccine to be effective.
- Maximum protection is typically reached a few weeks after your vaccination.



Common Side Effects

- Injection site soreness
- Fatigue
- Headache
- Muscle aches
- Nausea

Side effects typically last 1-2 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519

You received the Johnson & Johnson vaccine.

- You need **1 dose** for this vaccine to be effective.
- Maximum protection is typically reached a few weeks after your vaccination.



Common Side Effects

- Injection site soreness
- Fatigue
- Headache
- Muscle aches
- Nausea

Side effects typically last 1-2 days. If side effects last longer than 3 days, call 614-645-1519 to get a free COVID-19 test.

Side Effect Relief

- If you have pain or discomfort, talk to your health care provider about taking an over-the-counter medicine such as ibuprofen or acetaminophen.
- If you have pain or discomfort where you got the shot:
 - Apply a clean, cool, wet washcloth over the area.
 - Use or exercise your arm.
- If you have a fever:
 - Drink plenty of fluids.
 - Dress lightly.

Questions?

- www.columbus.gov/coronavirus
- 614-645-1519