

Columbus Public Health - Ben Franklin Tuberculosis Control Program

2011 TB Risk Assessment for Franklin County, Ohio

DATA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
TB Cases in Franklin County	75	59	61	55	77	85	78	61	41	66
Case rates per 100,000 in Franklin County	6.9	5.4	5.6	5.1	7.1	7.8	7.2	5.5	3.63	5.85
TB Cases for Ohio:	306	257	229	219	260	240	252	213	180	182
Case rate per 100,000 for Ohio:	2.7	2.3	2.0	1.9	2.3	2.1	2.2	1.9	1.6	1.6

- ✓ In 2010 Franklin County, with 66 counted cases and (1) transfer in TB case, had the highest incidence of active TB in the state of Ohio (case rate, 5.85/100,000), caring for 36% of all Ohio's TB cases. Cuyahoga County had 36 TB cases with a case rate of 2.8 per 100,000, and Hamilton County had 27 cases with a case rate of 3.2 per 100,000 population.
- ✓ Of Franklin County's 2010 active TB cases, 60% came from countries with high incidences of TB. Because of the high INH resistance rate in this county, four-drug therapy should be initiated for all active TB cases in Franklin County.
- ✓ If you are following a patient for latent Tuberculosis infection (LTBI) the recommended treatment is Isoniazid (INH) for 9 months. LTBI usually is diagnosed by a positive skin test/ QuantiFERON TB-Gold Blood Test and a chest x-ray showing no evidence of active disease. LTBI is not a reportable disease.
- ✓ **For information, hours, directions, or to leave a message call 645-2199 and your call will be returned. Report active and suspect TB cases, as well as all positive AFB smear results, to Kim Wilkins, TB Registrar, at 645-1823 (fax, 645-8669). We must be involved in the case management of active TB patients that are being treated by private providers.**

Ben Franklin TB Control Program, 240 Parsons Avenue, Columbus, Ohio, 43215, www.publichealth.columbus.gov